

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

SECTION 24(C) REGISTRATION FOR SPECIAL LOCAL NEED FOR SALE AND USE ONLY IN THE STATE OF NORTH DAKOTA



MUSTANG MAX™ EC Insecticide

All Applicable Directions, Restrictions and Precautions on the Registered Product Label for Mustang Max EC Insecticide (EPA Registration Number 279-3327 are to be followed.

These directions for use must be in the possession of the user at the time of application.

EPA SLN Number: ND-080002

Expires September 1, 2009

CROP	PEST		USE RATE
FLAX	Aphids Cutworms Diamondback Moth Loopers Lepidopterous Larvae Flea Beetle Fleahoppers	Grasshopper Plant Bug Stink Bugs Seedpod Weevil Thrips Whitefly Armyworms	4.0 oz/A (0.025 lb ai/A)

DIRECTIONS FOR USE:

Apply Mustang Max EC at the rate of 0.025 lbs ai/ac (4.0 ounces/acre)

Make applications when insect populations reach economic thresholds. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results.

Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage (minimum of 3 gallons per acre by air and 10 gallons per acre by ground).

RESTRICTIONS:

Observe all spray drift precautions on the EPA registered label.

Do not apply within 7 days of harvest of flax.

Do not make applications less than seven days apart.

Do not apply more than 0.15 pound active ingredient per acre per season of Mustang Max EC.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG MAX EC LABEL (EPA Registration No 279-3327) MUST BE FOLLOWED.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia PA 19103
MustangMaxEC_24c_ND_Flax_07-18-2008

Mustang and -FMC Trademarks
©2008 FMC Corporation