

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Durango™

EPA Reg. No. 62719-517

**24(c) Special Local Need Registration SLN ND-050008
(For Distribution and Use Only in the State of North Dakota)**

Pre-Harvest Application in Flax

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Durango™ herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Durango according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Durango.

Directions for Use

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

Refer to the label booklet for Durango for general information on product use, mixing, application equipment and techniques, labeled uses, and weed rate tables.

Durango may be applied as a spot treatment or a broadcast application to mature flax infested with Canada thistle or perennial sowthistle. Ground or aerial broadcast applications may be made. Apply when the crop is physiologically mature (contains 30% or less grain moisture) and nearly ready to harvest.

Broadcast Application

Durango may be applied as an over-the-top broadcast spray to control labeled weeds. Apply up to 3/4 quart in a spray volume of 3 to 20 gallons of water per acre.

Spot Application

A 2% solution is recommended when making spot applications using hand-held or backpack sprayers. Spray coverage should be uniform and complete, but do not spray weed foliage to the point of runoff.

Specific Use Precautions:

Greater efficacy against perennial weeds may be achieved if a period of 10 to 14 days is allowed between treatment and crop harvest. Do not apply when heavy rainfall is anticipated or when irrigation is scheduled soon after application. To minimize drift, use nozzles that emit medium to large-sized droplets. The likelihood of drift from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. Extreme care must be used when making preharvest applications to avoid injury to desirable plants and crops.

Specific Use Restrictions:

Apply at least 7 days before harvest. Only one application per year may be made. Do not feed flax straw or fodder from treated crop to livestock. Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the label for Durango. Do not add surfactants, additives containing surfactant, buffering agents, or pH adjusting agents to spray mixtures of Durango applied preharvest.

Chemigation: Do not apply this product through any type of irrigation system.

Expiration Date: June 28, 2010.

TMTrademark of Dow AgroSciences LLC

R316-020

Accepted 06/28/05

Initial printing.