

AgriSOLUTIONSTM



2,4-D LV4

PULL HERE TO OPEN ▲


ACTIVE INGREDIENT:

2-ethylhexyl ester of 2,4-dichlorophenoxyacetic acid* 66.2%
OTHER INGREDIENTS** 33.8%
TOTAL 100.0%

*Isomer Specific by AOAC Method No. 6.275 (13th Edition) 1980.
 **2,4-Dichlorophenoxyacetic acid equivalent 44.0%.

Contains petroleum distillates.

Contains 3.8 lbs. of 2,4-Dichlorophenoxyacetic acid per gallon.



KEEP OUT OF REACH OF CHILDREN

CAUTION

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Do not give **ANY** liquid to the person. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES: Hold eyelids open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Contains petroleum distillate – vomiting may cause aspiration hazard.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of medical emergency call toll free 1-877-424-7452.

EPA Reg. No. 1381-102

EPA Est. No. 70989-IA-001

0/0513/8

Distributed by

Winfield Solutions, LLC

P.O. Box 64589, St. Paul, MN 55164-0589

NET CONTENTS

2.5 GALLONS

(9.46 L)

PF-31080-1

PROOF

THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors, Unwind, and Dieline.**

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

PLEASE NOTE: Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors. **THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.**

WE CANNOT PROCESS THIS ORDER WITHOUT AN AUTHORIZED SIGNATURE

ARTWORK IS APPROVED

REVISED PROOF NEEDED

Signed _____

Date _____

Please Return To: _____

Please indicate the correct unwind position

| | | Unwind Positions | | | |
|---------------|---|------------------|----------|-----------|----------|
| | | Head-Off | Foot-Off | Right-Off | Left-Off |
| Wound Outside | 1 | | 2 | 3 | 4 |
| | | HEAD | FOOT | RIGHT | LEFT |
| Wound Inside | 5 | | 6 | 7 | 8 |
| | | HEAD | FOOT | RIGHT | LEFT |



4201 North Westport Avenue, Sioux Falls, South Dakota 57107
 Phone: (605) 978-0451 • Fax: (605) 978-0463

PROOF DATE: June 26, 2008

CUSTOMER: OMNIUM/WINFIELD

JOB NUMBER: 35321

LABEL SIZE: 6.75" x 6.75"

LEAFLET FLAT SIZE: 6.75" x 29.6875"

LEAFLET FOLDED SIZE: 6.75" x 6.0"

LABEL COLORS: BLACK

PATTERN VARNISH

LEAFLET "IN" COLORS: BLACK

LEAFLET "OUT" COLORS: BLACK PMS 640

dieline does not print

I want more options, fol-

ills or equipment, or oth-
centrate

Detergent and hot water.

the contents of this con-
he mechanical system is
is [40 CFR 170.240(d)(4-

gricultural pesticides [40
rents listed in the Worker
reduced or modified as

ossible, wash thoroughly

er is present, or to inter-
to treated areas. Drift or
rinsate.

chemical in areas where
around a cistern or well

ing/loading and disposal
dwater supplies. Use of
oading equipment on an

es, ornamentals, vegeta-
tities of the spray, which

in a way that will contact
n. For any requirements

This Standard contains
gricultural pesticides.
pecific instructions and
y interval. The require-

(continued)

AGRICULTURAL USE REQUIREMENTS (cont.)

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a secure area, in original container only. Do not store near feed or foodstuffs.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Plastic containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Metal containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300.

WEED LIST

2,4-D LV4 will control these plants and other 2,4-D susceptible species:

ANNUAL AND BIENNIAL WEEDS

Annual fanweed (field pennycress), annual yellow sweet clover, *beggarticks, bull thistle, burdock, carpetweed, chickweed, cocklebur, coffee-weed, common mullein, common evening primrose, cornflower, croton, galinsoga, goatsbeard, hemp, henbit, horseweed (maretail), jewelweed, jimsonweed, *knotweed, *kochia, lambsquarters, mallow (Venice, dwarf, little), marshelder, morning glory (common, ivy, woolly), musk thistle, mustards (except blue), pennycress, pepperweed (field), **pigweeds, poorjoe (woolly plantain), *prickly lettuce, puncturevine, purslane, ragweed (common and giant), rough fleabane, Russian thistle, salsify, shepherdspurse, stinkweed, *smartweeds (annual), sowthistle (annual or spiny), sunflower tansymustard, tumbleweed, velvetleaf, vetches, water primrose, *wild carrot, wild lettuce, wild parsnips, wild radish, wild sweet potato.

PERENNIAL WEEDS

*Alfalfa, *bindweeds (hedge, field and European), blue lettuce, *broom snakeweed, buckhorn plantain, buttercup, *Canada thistle, catnip, chamise, chicory, climbing milkweed, curly indigo, dandelion, *docks, *dogbanes, *goldenrod, *ground ivy, *hawkweed (orange), *hoary cress, *Jerusalem artichoke, locoweed, *many-flowered aster, milkvetch, *netties, nutgrass, plantains, poison ivy, pokeweed, sheep sorrel, sicklepod, sneezeweed (bitter), sowthistle (perennial), *tansy ragwort, *vervains, *wild garlic, wild onion, witchweed, wormwood, yellow rocket, yellow starthistle.

*BRUSH

Boxelder, buckbrush, coyotebrush, elderberry, manzanita, rabbitbrush, sagebrush (coastal, big, sand), sand shinnery oak, sumac, willow.

*These species may require repeat treatments and/or the specified higher rate.

**Control of pigweeds in the Texas and Oklahoma High Plains may be difficult. Do not exceed maximum application rates specified.

USE DIRECTIONS

Do not apply this product through any type of irrigation system.

Unless noted otherwise under individual **DIRECTIONS** section, for aerial application, apply the specified amount in a minimum of 2 gallons of water per acre. For ground application, apply the specified amount in a minimum of 3 gallons of water per acre. Use more water for both methods when adverse growing conditions are present. **DO NOT** apply with high spray pressures, hollow cone or other nozzle types that produce small spray droplets which may drift. Avoid spray drift by making applications when conditions such as wind, air stability and

temperature inversions a when weather is warm; expected within an hour. temperatures below free-

A variety of factors inclu cation (e.g., ground, aeri when applying this prod 2,4-D esters may volatiliz temperatures.

Droplet Size
When applying sprays th ents that require a coars microns or greater for sp When applying sprays th or coarser spray (ASAE s

Wind Speed
Do not apply at wind spe sensitive areas (including 250 feet downwind. If ap

Temperature Inversio
If applying at wind spec a) conditions of tempera b) stable atmospheric c atmospheric conditio

Susceptible Plants
Do not apply under circ rendered unfit for sale, u fruit trees (foliage), soyb of spray drift that might i

Other State and Local
Applicators must follow a gent regulations, they m


Equipment
All aerial and ground app *Additional requirements* : The boom length must r with efficacy and flight si for aircraft safety. This re the swath will be displac *Additional requirements* : Do not apply with a nozz

WATER BASED SPRAY WATER AND SOYBEAN

SOYBEAN OIL OR PET
mixture, if it does, an inv

NITROGEN FERTILIZE
use rates. Add half the fe ish filling the spray tank is a problem, the use of t recommendations of **Agri** of fertilizer use. Lower us

Adjuvants for Preeme
be added to the spray sc



temperature inversions are not a factor. The use of a suitable drift control agent at the proper rate will aid in the reduction of spray drift. Apply when weather is warm and plants are rapidly growing. Cold weather or dry conditions may cause poor results. **DO NOT** apply if rain is expected within an hour. Consult your local agronomist or Extension specialist for specific use and crop tolerance situations. When stored at temperatures below freezing, it may be necessary to warm contents to 45° F. and mix thoroughly before using.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if:

- a) conditions of temperature inversion exist, or
- b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications. When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

MIXING INSTRUCTIONS

WATER BASED SPRAY – Fill the equipment half full of water, agitate while adding this product, then add the rest of water.

WATER AND SOYBEAN OIL OR PETROLEUM OIL-BASED SPRAY – First mix this product with the oil, then add to water. If vigorous agitation is possible, the oil can be added last. **DO NOT ADD OIL FIRST!**

SOYBEAN OIL OR PETROLEUM OIL-BASED SPRAY: Add this product to straight oil to form a solution. Do not allow water to get into this mixture, if it does, an invert emulsion will form.

NITROGEN FERTILIZER: Weed and feed applications for corn, small grains, grasses grown for seed or grass pastures according to label use rates. Add half the fertilizer to the tank, then add specified label amount of **2,4-D LV4** per acre. Agitate constantly and vigorously and finish filling the spray tank with fertilizer. Apply as soon as possible, agitating constantly. Do not hold spray mixture overnight. If incompatibility is a problem, the use of **COMPLETE COMPATIBILITY™** agent at the specified label rate may correct the problem. Fertilize according to the recommendations of **Agri-Source™** labs, your supplier, or your Extension specialist. Herbicide foliage contact burning may occur as a result of fertilizer use. Lower use rates and concentrations will reduce this problem.

Adjuvants for Preemergence and Preplant Applications: A non-ionic surfactant such as **PREFERENCE®** or a crop oil concentrate may be added to the spray solution when this product is applied preemergence or preplant to increase control of large or difficult to control weeds.

involves contact with any-

Standard for agricultural farms, forests, nurseries, or

or rinsate is a violation of instructions, contact your Regional Office for guidance. rilling, or puncture and dis- stay out of smoke. Metal sanitary landfill, or by other

-800-424-9300.

chickweed, cocklebur, co- horseweed (marestalk), jew- common, ivy, woolly), musk ice, puncturevine, purslane, (annual), sowthistle (annual) d parsnips, wild radish, wild

up, *Canada thistle, catnip, weed (orange), *hoary cress, red, sheep sorrel, sicklepod, wood, yellow rocket, yellow

ry oak, sumac, willow.


on rates specified.

t in a minimum of 2 gallons re. Use more water for both o or other nozzle types that h as wind, air stability and

Crop oil concentrates tion). Non-ionic surfac Wash spray equipment the ground: spray or Apply the specified a

CORN – All cor

| WEED |
|--|
| Preplant – Annual a broadleaf seedlings Perennial weed seed and existing cover c |
| *Do not perform tillage |
| Preemergence and tillage – Broadleaf w and annual grasses |
| *Do not use on sanc |
| Postemergence – A broadleaf weeds |
| Perennial broadleaf |
| **DO NOT apply frc oil or other adjuvant a week to 10 days a |
| Late season weed c Preharvest (Field co popcorn only) |
| *Use lower rate for s |
| RESTRICTIONS A |
| <ul style="list-style-type: none"> • Do not use treat • The preharvest ir • Maximum of 5.2 • Preplant or preer Limited to one pi Maximum of 2.1 • Postemergence: Limited to one pi Maximum of 1 pt • Preharvest: Limited to one pi Do not exceed 2 |
| RESTRICTIONS A |
| <ul style="list-style-type: none"> • Do not use treat • The preharvest ir • Minimum of 21 d • Maximum of 3.15 • Preplant or preer Limited to one pi Maximum of 2.1 • Postemergence: Limited to one pi Maximum of 1 pt |



reduction of spray drift. Apply suits. **DO NOT** apply if rain is in sight. When stored at 100°F (38°C) or higher, the product may become viscous. Do not use if the product is thick or clumpy.

humidity) and method of application. Make appropriate adjustments.

conditions of low humidity and high temperatures.

2,4-D mixed with active ingredients. The maximum volume mean diameter of 385 microns.

When spraying, apply only as a medium spray using medium sized nozzles.

Do not apply if there are not any weeds (weeds, non-target crops) within the treated field.

Temperature inversions or stable conditions.

Do not be damaged or crops thereof (peaches, grapes (in growing stage), apples, pears, or tobacco. Small amounts of spray may be applied to these crops.

Where states have more stringent restrictions, use those restrictions.

Use other surrogates.

Apply at the lowest height consistent with the label. If a greater height is required, the nozzle is made with a crosswind, aircraft upwind.

Do not rest of water. Do not add to water. If vigorous agitation is required, use a mechanical agitator.

Do not allow water to get into this product.

Do not use in pastures according to label directions. Do not apply to pastures and finish overnight. If incompatibility occurs, fertilize according to the label. Fertilizing may occur as a result of application.

Do not use a crop oil concentrate may be used or difficult to control weeds.

Crop oil concentrates must contain at least 17% emulsifier, and should be used at 1% volume/volume (1 gallon per 100 gallons of spray solution). Non-ionic surfactants should be used at a 0.25% volume/volume (1 quart per 100 gallons of spray solution).

Wash spray equipment thoroughly with **PROTANK™** cleaner after using this product. When cleaning equipment, do not pour washwater on the ground: spray or drain over a large area away from wells or other water sources.

Apply the specified amount of 2,4-D per acre regardless of the amount of diluent used.

CORN – All corn including FIELD, SWEET and POP

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|--|------------------------------|---|
| Preplant – Annual and biennial broadleaf seedlings | 1 pint | Planting of corn must be delayed a minimum of 7 days after application at rates up to 1 pint per acre, and a minimum of 14 days at rates from 1 to 2 pints per acre. Planting sooner after application than specified on this label may result in unacceptable crop injury. |
| Perennial weed seedlings and existing cover crops | 1 to 2.1 pints* | |
| *Do not perform tillage for at least 7 days after application. Do not use on sandy soils or unacceptable crop injury may result. | | |
| Preemergence and reduced tillage – Broadleaf weeds and annual grasses | 2.1 pints* | Apply after corn is planted but before emergence for control of emerged broadleaf weeds. The seed furrow must be completely closed at application or severe crop injury may result. |
| *Do not use on sandy soils or unacceptable crop injury may result. | | |
| Postemergence – Annual broadleaf weeds | 1/2 pint | Apply when corn is less than 8 inches tall, but to avoid crop injury do not apply just after leaves have unfolded. If corn is over 8 inches tall, use drop nozzles to keep spray off of corn foliage as much as possible. See additional restrictions below. |
| Perennial broadleaf weeds | 3/4 to 1 pint** | Apply when weeds are in bud to bloom stage. If corn is over 8 inches tall, use drop nozzles to keep spray off corn foliage as much as possible. Sweet Corn: To minimize potential for crop injury, use only lowest rate in rate range. |
| **DO NOT apply from 2 weeks before tasseling to dough stage. DO NOT apply to open whorls. To avoid injury, do not use with atrazine, oil or other adjuvants. Application during high moisture and temperature conditions may cause injury or brittleness. DO NOT cultivate for a week to 10 days after treatment or stalk breakage may occur. | | |
| Late season weed control – Preharvest (Field corn and popcorn only) | 1 to 2 pints* | Apply after silks are completely brown (denting stage) to reduce weeds that interfere with harvest and to decrease weed seed production. Do not apply to sweet corn. |
| *Use lower rate for small annual and biennial weeds. Use the higher rate for perennial and larger hard-to-kill annual and biennial weeds. | | |
| RESTRICTIONS AND LIMITATIONS FOR FIELD CORN AND POPCORN: | | |
| <ul style="list-style-type: none"> Do not use treated crop as fodder for 7 days following application. The preharvest interval (PHI) is 7 days. Maximum of 5.2 pts. (2.42 lbs. ae) per acre per crop cycle. Preplant or preemergence: Limited to one preplant or preemergence application per crop cycle. Maximum of 2.1 pts. (1 lb. ae) per acre per application. Postemergence: Limited to one postemergence application per crop cycle. Maximum of 1 pt. (1/2 lb. ae) per acre per application. Preharvest: Limited to one preharvest application per crop cycle. Do not exceed 2 pts. per acre per application for preharvest use. | | |
| RESTRICTIONS AND LIMITATIONS FOR SWEET CORN: | | |
| <ul style="list-style-type: none"> Do not use treated crop as fodder for 7 days following application. The preharvest interval (PHI) is 45 days. Minimum of 21 days between applications. Maximum of 3.15 pts. (1-1/2 lbs. ae) per acre per crop cycle. Preplant or preemergence: Limited to one preplant or preemergence application per crop cycle. Maximum of 2.1 pts. (1 lb. ae) per acre per application. Postemergence: Limited to one postemergence application per crop cycle. Maximum of 1 pt. (1/2 lb. ae) per acre per application. | | |

SOYBEANS

| WEEDS |
|--|
| Preplant – Emergent broadleaf weeds |
| USE RESTRICTIONS: sandy soils or unsuitable for 2,4-D preplant: <ul style="list-style-type: none"> The maximum of 1 pint per acre. Preplant: Limited to 1 pint per acre. Maximum of 1 pint per acre. Apply not less than 14 days before planting. OR Preplant: Limited to 1 quart per acre. Maximum of 2 quarts per acre. Apply not less than 14 days before planting. |

GRAIN SORGHUM

| WEEDS |
|--|
| Annual broadleaf |
| Perennial broadleaf |
| RESTRICTIONS: <ul style="list-style-type: none"> The preharvest interval (PHI) is 7 days. Do not permit application to grain. Postemergence: Limited to 1 quart per acre. Maximum of 1 quart per acre. |

flon per 100 gallons of spray solution).
 ment, do not pour washwater on

IS

um of 7 days after application
 imum of 14 days at rates from
 pplication than specified on this
 r.

op injury may result.

ergence for control of emerged
 e completely closed at applica-

ill, but to avoid crop injury do
 if corn is over 8 inches tall, use
 iage as much as possible. See

ige. If corn is over 8 inches tall,
 iage as much as possible.
 p injury, use only lowest rate in

injury, do not use with atrazine,
 rittleness. **DO NOT** cultivate for

ing stage) to reduce weeds that
 d seed production. Do not apply


kill annual and biennial weeds.

SOYBEANS

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|---|------------------------------|--|
| Preplant – Emerged broadleaf weeds | 1 to 2.1 pints* | After applying, plant soybean seed as deep as practical or at least 1-1/2 to 2 inches deep. Seed furrow must be completely closed or severe crop injury will result. |
| <p>USE RESTRICTIONS AND LIMITATIONS FOR SOYBEANS: Do not perform tillage for at least 7 days after application. Do not use on sandy soils or unacceptable crop injury may result. Do not replant treated fields in the same growing season with crops that are not labeled for 2,4-D preplant use.</p> <ul style="list-style-type: none"> The maximum rate per crop cycle is 2.1 pints (1 lbs. ae) per acre. Preplant: <ul style="list-style-type: none"> Limited to 2 preplant applications per crop cycle. Maximum of 1 pint (1/2 lb. ae) per acre per preplant application. Apply not less than 7 days prior to planting soybeans. <p>OR</p> <ul style="list-style-type: none"> Preplant: <ul style="list-style-type: none"> Limited to 1 application per crop cycle. Maximum of 2.1 pints (1 lb. ae) per preplant application. Apply not less than 15 days prior to planting soybeans. | | |

GRAIN SORGHUM

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|--|------------------------------|---|
| Annual broadleaf weeds | 1/2 pint | Apply to plants that are 5 to 15 inches tall. DO NOT treat plants less than 5 inches tall or from boot to early dough stage. Use drop nozzles when crop is 8 inches or taller. |
| Perennial broadleaf weeds | 2/3 to 1 pint | The higher rate may be needed for some weeds, but chances of crop injury may increase. DO NOT use oil. Some varieties and hybrids are 2,4-D sensitive. Crop injury may also be increased by high moisture and temperature conditions. Check with your seed company and Extension Service for advice. |
| <p>RESTRICTIONS AND LIMITATIONS FOR GRAIN SORGHUM:</p> <ul style="list-style-type: none"> The preharvest interval (PHI) is 30 days. Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application. Postemergence: <ul style="list-style-type: none"> Limited to 1 application per crop cycle. Maximum of 1 pt. (1/2 lb. ae) per acre per application. | | |



SMALL GRAINS – (Not underseeded with legumes)

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|--|--|--|
| Postemergence – Spring wheat, barley, and rye – Annual and biennial weeds Perennial broadleaf weeds | 1/2 to 2 pints* 1 to 2 pints** | Apply when grain is in full tiller stage (4 to 8 inches high) but before boot stage (Zadoks 2 ₅ to 4 ₀) when weeds are small and actively growing. Up to 2 pints per acre may be used to control difficult weed problems, but do not use unless some crop damage is acceptable. Apply only in the spring when crop is fully tillered, but before grain is in boot stage (before Zadoks 4 ₀). For improved control of difficult weeds, apply up to 2 pints per acre. |
| **DO NOT USE THE HIGHER RATE IF POSSIBLE CROP INJURY IS NOT ACCEPTABLE | | |
| Spring and winter wheat and barley Resistant weeds | 2,4-D LV4 may be used in combination with Ally™, Harmony Extra™, Express™, Finesse™, at their earlier application intervals to control resistant weeds such as kochia and Russian thistle. Follow application directions on each product label. | |
| Winter wheat and rye Annual weeds | 1/2 to 1 pint** | Apply only in the spring when crop is fully tillered, but before grain is in boot stage (before Zadoks 4 ₀). For improved control of difficult weeds, apply up to 2 pints per acre. |
| **DO NOT USE THE HIGHER RATE IF POSSIBLE CROP INJURY IS NOT ACCEPTABLE | | |
| Wild garlic or onions | 1/2 to 2 pints* | Apply 1-1/2 pints when grain is at full tiller and wild garlic and onion plants are small. Apply 2 pints after harvest to the crop stubble. For control of new fall growth of these plants, refer to fallow land use directions. |
| Spring-seeded oats Fall seeded oats grown for grain (Southern) | 1/2 to 2/3 pint* 2/3 to 1-1/2 pints* | Apply at full tiller, but before early boot stage (Zadoks 2 ₅ to 4 ₀). Apply at full tiller, but before early boot stage (Zadoks 2 ₅ to 4 ₀). |
| *Difficult-to-control weeds may require higher rate, but some injury may occur since oats are less tolerant to 2,4-D than wheat or barley. DO NOT spray during or just after cold weather. | | |
| Preharvest – Cereal grains | 1 pint | Apply when grain is in hard dough stage (Zadoks 8 ₇) to control weeds that will interfere with harvest. Apply when soil moisture is adequate for weed growth for best results. |
| *Use the lower rate for small annual and biennial weeds. Use the higher rate for perennial weeds or hard-to-kill annual or biennial weeds. The higher rate should be used only where heavy weed infestation is a problem and increased risk of crop damage is acceptable. | | |
| RESTRICTIONS AND LIMITATIONS FOR SMALL GRAINS: | | |
| <ul style="list-style-type: none"> The preharvest interval (PHI) is 14 days. Postemergence: Limited to one postemergence application per crop cycle. Maximum of 2 pts. (1 lb. ae) per acre per application. Preharvest: Limited to one preharvest application per crop cycle. Maximum of 1 pt. (1/2 lbs. ae) per acre per application. Limited to 3 pts. (1.425 lbs. ae) per acre per crop cycle. | | |

FALLOW GROUND (cr

| WEEDS |
|--|
| Annual broadleaf weeds |
| Biennial weeds |
| Perennial weeds |
| Wild onions and garlic |
| RESTRICTIONS AND LIMITATIONS: |
| <ul style="list-style-type: none"> Plant only labeled crops Limited to 2 applications Maximum of 4.2 pts. (2 lb) Minimum of 30 days betw |
| PLANTING IN TREATED AREAS: |
| <p>Labeled Crops: Within 29 c</p> <p>Follow more specific limitatio soon after application, espec</p> <p>Other Crops: All other crop However, under certain cond sidered in weighing this risk. tures above freezing have eli</p> <p>Degradation Factors: Whe tions following application he product were applied and sc tion. Consult your local agric</p> |
| CONSERVATION RESTRICTIONS: |
| WEEDS |
| Annual broadleaf and perennial weeds |
| **Buckbrush, coyotebrush, rabbitbrush, sagebrush, and other chaparral species |
| **Sand shinnery oak |
| *Deep-rooted perennial weed |
| **Woody plants and any regre |
| RESTRICTIONS AND LIMITATIONS FOR CULTURAL PRODUCTION: |
| <ul style="list-style-type: none"> The preharvest interval (PHI) is 14 days. Postemergence: Limited to 2 applications Maximum of 4.2 pts (2 lbs) Minimum of 30 days betw If grass is to be cut for ha For program lands, such The more restrictive requi |

FALLOW GROUND (crop stubble on idle land, or postharvest to crops, or between crops)

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|------------------------|------------------------------|---|
| Annual broadleaf weeds | 1 to 2 pints | Use the lower rate for small actively growing weeds. Use the higher rate on larger or weather stressed weeds. |
| Biennial weeds | 2 to 4.2 pints | Use the lower rate in the spring on biennial weeds such as the musk thistle during the rosette stage before stalks have formed. Use the higher rate after stalk formation or in the fall. |
| Perennial weeds | 2 to 4.2 pints | Apply during the bud to bloom stage while weeds are actively growing. Do not till for 2 weeks after treatment or until the weeds start to die. |
| Wild onions and garlic | 4.2 pints | Apply to regrowth in fall after harvest. |

RESTRICTIONS AND LIMITATIONS FOR FALLOW GROUND crop stubble on idle land, or postharvest to crops, or between crops:

- Plant only labeled crops within 29 days following application.
- Limited to 2 applications per year.
- Maximum of 4.2 pts. (2 lbs. ae) per acre per application.
- Minimum of 30 days between applications.

PLANTING IN TREATED AREAS:

Labeled Crops: Within 29 days after an application of this product, plant only those crops listed on this or other registered 2,4-D labels. Follow more specific limitations, if any, provided in directions for specific crops. Labeled crops may be at risk of crop injury or loss if planted soon after application, especially during the first 14 days. Degradation factors described below should be considered in weighing this risk.

Other Crops: All other crops may be planted 30 or more days after application without concern for illegal residues in the planted crop. However, under certain conditions, there may be a risk of injury to susceptible crops. Degradation factors described below should be considered in weighing this risk. Under normal conditions, any crop may be planted without risk of injury if at least 90 days of soil temperatures above freezing have elapsed since application.

Degradation Factors: When planting into treated areas, the risk of crop injury is less if lower rates of product were applied and conditions following application have included warm, moist soil conditions that favor rapid breakdown of 2,4-D. Risk is greater if higher rates of product were applied and soil temperatures have been cold and/or soils have been excessively wet or dry in the days following application. Consult your local agricultural extension service or information about susceptible crops and typical conditions in your area.

CONSERVATION RESERVE PROGRAM AREA PASTURES AND RANGELAND NOT IN AGRICULTURAL PRODUCTION

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|---|------------------------------|---|
| Annual broadleaf and *perennial weeds | 3 pints | Do not apply after heads form or when grass is in boot to milk stage when a seed crop is desired. DO NOT use on alfalfa, clover, other legumes, or newly seeded pastures. For aerial application, apply the recommended amount in a minimum of 2 gallons of water per acre. For ground application, use a minimum of 10 gallons of water per acre. |
| **Buckbrush, coyotebrush, rabbitbrush, sagebrush, and other chaparral species | 4.2 pints | Apply in 5 to 10 gallons of water plus 1 gallon of oil or 1-2 quarts of a crop oil concentrate with at least 17% emulsifiers per acre or a non-ionic surfactant at .25% v/v surfactant to water – (1 quart per 100 gallons of water) per acre. |
| **Sand shinnery oak | 4.2 pints | Apply 5 gallons of oil or 4 gallons of water plus 1 gallon of oil or 1-2 quarts of a crop oil concentrate with at least 17% emulsifiers per acre or a non-ionic surfactant at .25% v/v surfactant to water – (1 quart per 100 gallons of water) per acre. |

*Deep-rooted perennial weeds may require the higher rate or repeated treatments.
 **Woody plants and any regrowth may require repeat treatments.

RESTRICTIONS AND LIMITATIONS FOR CONSERVATION RESERVE AREA PASTURES AND RANGELAND NOT IN AGRICULTURAL PRODUCTION:

- The preharvest interval (PHI) is 7 days (cut forage for hay).
- Postemergence:
 Limited to 2 applications per year.
 Maximum of 4.2 pts (2 lbs. ae) per acre per application.
 Minimum of 30 days between applications.
 If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.
 For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used.
 The more restrictive requirements of the program rules or this label must be followed.

PASTURES AND RA cultural production, ex

For susceptible annual and biennial weeds:
 For moderately susceptible perennial weeds:
 For spot treatment: Do not apply more than 4.2 pts (2 lbs. ae) per acre.

RESTRICTIONS AND LI

- Do not cut forage for h
- Maximum of two applic
- Do not apply more than
- The minimum retreatme

NON-CROP AREAS and utility rights-of-w

WEEDS

- Annual broadleaf weeds
- Perennial and biennial broadleaf weeds
- Tansy ragwort and musk t
- Wild onion and wild garlic
- Woody plants – Ground application

- Woody plants – Aerial application
- Spot treatment

RESTRICTIONS AND LI

- Postemergence (annual and biennial weeds):
 Limited to 2 applicator
 Maximum of 4.2 pts. (2 lbs. ae) per acre per application.
- Postemergence (woody plants):
 Limited to 1 applicator
 Maximum of 6 pts. (2.8 lbs. ae) per acre per application.
- Application to non-crop areas: For commercial use, or for

TURF, ORNAMENTAL

WEEDS

- Annual broadleaf weeds
- Biennial and perennial weeds
- *Deep-rooted perennials n

RESTRICTIONS AND LI and other lawn and gra

- Postemergence:
 Limited to 2 applicator
 Maximum of 3.15 pts. (1.575 lbs. ae) per acre per application.
 The maximum seasona

es high) but before and actively growing. weed problems, but a. Apply only in the n boot stage (before pply up to 2 pints per

M, Finesse™, at Russian thistle.

before grain is l of difficult weeds,


glic and onion plants le. For control of : directions.

2₅ to 4₀). 2₅ to 4₀).

han wheat or barley.

o control weeds that adequate for weed

al or biennial weeds. is acceptable.



1 crops)

the higher rate on
the musk thistle dur-
igher rate after stalk
vely growing. Do not
ie.
o crops, or between

registered 2,4-D labels.
injury or loss if planted
ad in weighing this risk.
ies in the planted crop.
J below should be con-
J days of soil tempera-
ere applied and condi-
reater if higher rates of
days following applica-
s in your area.

AGRICULTURAL

ot to milk stage when
ther legumes, or newly
nded amount in a min-
n, use a minimum of 10

-2 quarts of a crop oil
non-ionic surfactant at
water) per acre.

of oil or 1-2 quarts of a
or a non-ionic surfac-
ions of water) per acre.

AND NOT IN AGRI-

s or hay may be used.

PASTURES AND RANGELAND (established grass pastures, rangeland, and perennial grasslands not in agri-cultural production, excluding Conservation Reserve areas)

For susceptible annual and biennial broadleaf weeds: Do not apply more than 2.1 pts (1 lb. ae) per acre per application.
For moderately susceptible biennial and perennial broadleaf weeds and difficult-to-control weeds and woody plants: Do not apply more than 4.2 pts (2 lbs. ae) per acre per application.
For spot treatment: Do not apply more than 4.2 pts (2 lbs. ae) per acre.

- RESTRICTIONS AND LIMITATIONS FOR PASTURES AND RANGELAND:**
- Do not cut forage for hay within 7 days of application.
 - Maximum of two applications per year.
 - Do not apply more than 4 lbs. ae/acre per year.
 - The minimum retreatment interval is 30 days.

NON-CROP AREAS – Fencerows, hedgerows, roadsides, ditches, rights-of-way, airfields, railroads, highway and utility rights-of-way, industrial sites, and other non-crop areas

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|--|------------------------------|--|
| Annual broadleaf weeds | 2 to 4.2 pints | Apply when weeds are young and growing vigorously. |
| Perennial and biennial broadleaf weeds | 2 to 3 pints | Spray perennial weeds when near the bud stage, but not flowering. Do not use on St. Augustine grass. Bentgrass, clover, legumes and dichondra may be injured. Do not apply to newly seeded areas until grass is well established. Deep-rooted perennials may require repeated treatments. |
| Tansy ragwort and musk thistle | | Apply in rosette stage before bolting. |
| Wild onion and wild garlic | | Treat in the early spring and fall when young and actively growing. |
| Woody plants – Ground application | 6 pints | Apply in 20 to 100 gallons of water. For increased effectiveness, add a crop oil concentrate with at least 17% emulsifiers at 1-2 quarts per acre or a non-ionic surfactant at .25% v/v surfactant to water – 1 quart per 100 gallons of water. Spray volumes of up to 500 gallons per acre may be needed for control if brush is dense. |
| Woody plants – Aerial application | 4 to 6 pints | For solid stands of susceptible brush, apply in 3 to 12 gallons volume per acre. 2 to 4 quarts of fuel oil may be included in this mixture. |
| Spot treatment | 4 ounces | Apply in 3 gallons of water, mixing thoroughly, and spray to runoff. This high dosage rate may only be used where injury may be tolerated. |

- RESTRICTIONS AND LIMITATIONS FOR NON-CROP AREAS:**
- Postemergence (annual and perennial weeds): Limited to 2 applications per year. Maximum of 4.2 pts. (2 lbs. ae) per acre per application.
 - Minimum of 30 days between applications.
 - Postemergence (woody plants): Limited to 1 application per year. Maximum of 6 pts. (2.85 lbs. ae) per acre per year.
 - Application to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

TURF, ORNAMENTAL (golf courses, parks, cemeteries, sports fields, turfgrass, and other lawn and grass areas)

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|------------------------------|------------------------------|--|
| Annual broadleaf weeds | 2 pints | Do not apply to newly seeded areas until grass is well established. Where bentgrass predominates, make 2 applications of 1 pint per acre at 3-week intervals. Do not use on susceptible southern grasses such as St. Augustine, bentgrass and dichondra. |
| Biennial and perennial weeds | 2 to 3.15 pints* | |

*Deep-rooted perennials may require repeat applications. Clovers and legumes may be injured by this treatment.

- RESTRICTIONS AND LIMITATIONS FOR TURF, ORNAMENTAL (golf courses, parks, cemeteries, sports fields, turfgrass, and other lawn and grass areas):**
- Postemergence: Limited to 2 applications per year. Maximum of 3.15 pts. (1-1/2 lbs. ae) per acre per application. The maximum seasonal rate is 6.3 pts. (3 lbs. ae) per acre, excluding spot treatments.


TURF GROWN FO

| WEEDS |
|-------------------------|
| Annual broadleaf weed |
| Perennial and biennial |
| *Use only the low rate |
| RESTRICTIONS AND |
| • Limited to 2 applica |
| • Maximum of 3 pts. 1 |
| • Minimum of 21 days |

The directions for use of th
TO YOU ARE FURNISHED
OR REPRESENTATIONS O
WITH REGARD TO THE PE
ELIGIBILITY OF THE PRO
FECTIVENESS, MAY RESL
THE GOODS, OR THE M/
TURER OR SELLER AND /
BUYER, MANUFACTUREF
RESENTATION OR WARR

TO THE EXTENT CONSIS
OR CONSEQUENTIAL DA
ICATION, HANDLING, AND
BUYER OR TO CUSTOME
FROM OR BY REASON O
GOODS SOLD BY MANUF
BUYER'S OR USER'S EX
THE COST OF THE PROD
If you do not agree with or
it unopened to the Seller, z

Ally™, Harmony Extra™, E



grasslands not in agri-

lication.
plants: Do not apply more

ds, railroads, highway

usly.
but not flowering. Do not
jumes and dichondra may
until grass is well estab-
treatments.
actively growing.
effectiveness, add a crop
2 quarts per acre or a non-
1 quart per 100 gallons of
e may be needed for con-
to 12 gallons volume per
his mixture.
I spray to runoff. This high
be tolerated.
ng grown for sale or other

lawn and grass areas)

grass is well established.
ions of 1 pint per acre at
thern grasses such as St.
t.
ports fields, turfgrass,

TURF GROWN FOR SEED OR SOD - (DO NOT USE IN CALIFORNIA)

| WEEDS | AMOUNT OF 2,4-D LV4 PER ACRE | DIRECTIONS |
|---|------------------------------|--|
| Annual broadleaf weeds | 1 to 1-1/2 pints* | Apply to established stands before the seed head comes into the boot stage. Do not spray in boot stage of growth. For seedling grasses, apply in the spring after grass has tillered or has at least 5 leaves, but before boot stage. |
| Perennial and biennial weeds | 2 to 3 pints* | |
| *Use only the low rate on seedling grasses. | | |
| RESTRICTIONS AND LIMITATIONS FOR TURF GROWN FOR SEED OR SOD: | | |
| <ul style="list-style-type: none"> Limited to 2 applications per year. Maximum of 3 pts. (1.42 lbs. ae) per acre per application. Minimum of 21 days between applications. | | |

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS' CUSTOMERS, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Ally™, Harmony Extra™, Express™, and Finesse™ are registered trademarks of E.I. duPont de Nemours and Company.

ACTIVE INGREDIENT
2-ethylhexyl este
OTHER INGREDIENT
TOTAL

*Isome

Cont:

Harmful if swallowed. (C
repeated skin contact r

IF SWALLOWED: C
ing unless told to by .
IF IN EYES: Hold ey
then continue rinsing.
IF ON SKIN: Take of
treatment advice.
IF INHALED: Move p
if possible. Call a poi

Have the product cor
medical emergency c

EPA Reg. No. 138

Distributed by
Winfie
P.O. Box 6458

Job #35321

ad comes into the boot stage.
seedling grasses, apply in the
st 5 leaves, but before boot

AW, (1) THE GOODS DELIVERED
IO WARRANTIES, GUARANTEES,
JE, STATUTORY OR OTHERWISE,
ARTICULAR PURPOSE, USE, OR
NG BUT NOT LIMITED TO INEF-
S USED IN COMBINATION WITH
) THE CONTROL OF MANUFAC-
S AND AGREEMENTS BETWEEN
AUTHORITY TO MAKE ANY REP-

BLE FOR SPECIAL, INCIDENTAL,
SOLD, INCLUDING USE, APPLI-
Y WAY OF INDEMNIFICATION TO
NDS WHATSOEVER, RESULTING
TRUCTIONS FOR USE, OF THE
R, USER, OR ITS' CUSTOMERS,
OR DAMAGES NOT EXCEEDING

to not use the product, and return



2,4-D LV4

ACTIVE INGREDIENT:

2-ethylhexyl ester of 2,4-dichlorophenoxyacetic acid* 66.2%

OTHER INGREDIENTS**

33.8%

TOTAL

100.0%

*Isomer Specific by AOAC Method No. 6.275 (13th Edition) 1980.

**2,4-Dichlorophenoxyacetic acid equivalent 44.0%.

Contains petroleum distillates.

Contains 3.8 lbs. of 2,4-Dichlorophenoxyacetic acid per gallon.



(01) 0 0784742 00005 8

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Do not give **ANY** liquid to the person. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eyelids open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Contains petroleum distillate - vomiting may cause aspiration hazard.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of medical emergency call toll free 1-877-424-7452.

EPA Reg. No. 1381-102

EPA Est. No. 70989-IA-001

0/0513/8

Distributed by

Winfield Solutions, LLC

P.O. Box 64589, St. Paul, MN 55164-0589

NET CONTENTS

2.5 GALLONS

(9.46 L)