

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

EMERGENCY INFORMATION	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY: Transportation: CHEMTREC 1-800-424-9300 Other: AMVAC 1-323-264-3910	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING: May cause eye irritation. Harmful if swallowed. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
 Applicators and other handlers must wear:
 • Long-sleeved shirt and long pants
 • Waterproof gloves
 • Shoes plus socks
 • Protective eyewear

USER SAFETY REQUIREMENTS
 • Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentration. Do not reuse them.
 • Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS
 This pesticide is extremely toxic to fish and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

User Safety Recommendations	
Users should: <ul style="list-style-type: none"> Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS	
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: <ul style="list-style-type: none"> Coveralls Waterproof gloves Shoes plus socks Protective eyewear 	

GENERAL INFORMATION
 Apply AMBUSH 25W as shown in "Directions for Application". Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rates may be required under conditions of heavy infestation. Do not apply more than 2.0 lbs. active ingredient (a.i.) per acre per season. Crops in this label may be planted immediately after last treatment. Do not plant other crops within 60 days after last application. Any other crop may be planted if it is not harvested or grazed.
Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your Local or State Agricultural Authorities for details.
 If resistance to this product develops in your area, this product or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may

be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

CHEMIGATION
SPRINKLER IRRIGATION APPLICATION

Apply AMBUSH 25W Insecticide at rates and timing described elsewhere in this label. Apply with center pivot or continuous-move equipment distributing 1/2 acre-inch or less during treatment. If stationary (solid sets, handlines or wheel lines other than continuous-move) are used, AMBUSH 25W insecticide should be injected into no more than the last 20 to 30 minutes of the set. Do not apply when winds are greater than 10 to 15 mph to avoid drift or wind skips. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

USE PRECAUTIONS-SPRINKLER IRRIGATION APPLICATION

- Apply this product only through sprinkler (including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Do not apply through chemigation systems connected to public water systems.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AI/Acre (OZ./Acre)	REMARKS
COLE CROPS - Brassica Leafy Vegetables			
Broccoli	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.
Brussels Sprouts	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 to 10 days or as needed. Do not apply more than 1.0 lb. a.i. per acre per season. Do not apply within 1 day of harvest.
Cabbage	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 to 10 days or as needed. Do not apply more than 1.0 lb. a.i. per acre per season. Do not apply within 1 day of harvest.
Cauliflower	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.

(continued)

DIRECTIONS FOR APPLICATION (continued)			
CROP	TARGET PESTS	RATE LBS AI/Acre (OZ./Acre)	REMARKS
COLE CROPS - Brassica Leafy Vegetables (continued)			
Chinese Broccoli ^b (gai lan, white flowering)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 3 days of harvest.
Chinese Cabbage (Tight-heading varieties only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 to 10 days or as needed. Do not apply more than 1.0 lb. a.i. per acre per season. Do not apply within 1 day of harvest.
Cucurbit Vegetables ^c	Leafminers and Squash Bugs Cabbage Looper Cucumber Beetle (adults) Cutworms Melon Worm Pickleworm Plant Bugs (including Lygus and Stink Bugs) Rindworms and Squash Vine Borer	0.2 (12.8) 0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. Apply a minimum of 4 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 lbs. a.i. per acre per season. AMBUSH 25W may be applied up to harvest.

(continued)

DIRECTIONS FOR APPLICATION (continued)			
CROP	TARGET PESTS	RATE LBS AI/Acre (OZ./Acre)	REMARKS
FRUITING VEGETABLES			
Eggplant	Colorado Potato Beetle Vegetable Leafminer Cabbage Looper Diamondback Moth Fall Armyworm Beet Armyworm Corn Earworm Green Cloverworm Leafhoppers European Corn Borer	0.2 (12.8) 0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 2.0 lbs. a.i. per acre per season. Do not apply within 3 days of harvest.
Peppers (Bell)	Vegetable Leafminer Cabbage Looper Flea Beetle Pepper Weevil European Corn Borer	0.1-0.2 (6.4-12.8) 0.2 (12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.6 lbs. a.i. per acre per season. Do not apply within 3 days of harvest.
Tomatoes	Tomato Fruitworm Cabbage Looper Tomato Pinworm Vegetable Leafminer Beet Armyworm Southern Armyworm Colorado Potato Beetle Granulate Cutworm Hornworm	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Do not apply more than 1.2 lbs. a.i. per acre per season. Do not apply as a band. Do not use on tomato varieties in which the mature tomatoes will be less than 1 inch in diameter (such as cherry tomatoes).

(continued)

DIRECTIONS FOR APPLICATION (continued)			
CROP	TARGET PESTS	RATE LBS AI/Acre (OZ./Acre)	REMARKS
LEAFY VEGETABLES			
Leafy Vegetables ^a (except Brassica)	Vegetable Leafminer Cabbage Looper Cutworm Complex Fall Armyworm Beet Armyworm Corn Earworm Green Cloverworm Leafhoppers European Corn Borer	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply every 3 to 5 days or as needed. Do not apply more than 2.0 lbs. a.i. per acre per season. Can be applied up to 1 day of harvest.
Onions (Dry Bulb and Garlic)	Onion Thrips Armyworms Cutworms Leafminers Onion Maggots (adult) Stink Bugs	0.15-0.3 (9.6-19.2) 0.1-0.3 (6.4-19.2)	Apply by air or ground equipment. Apply in a minimum of 20 gallons of water per acre with ground equipment or in a minimum of 6 gallons of water per acre by air as foliar spray. Begin applications when pests appear and repeat as necessary to maintain control. Use the higher label rates as onion thrips population increases and avoid rescue situations. Do not apply more than 2.0 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.
Potatoes	Colorado Potato Beetle Potato Leafhopper Aster Leafhopper Potato Flea Beetle Potato Aphid Potato Tuberworm Potato Psyllid Cabbage Looper	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply as needed. Do not apply more than 1.6 lbs. a.i. per acre per season. Can be applied up to 14 days prior to harvest.

(continued)

DIRECTIONS FOR APPLICATION (continued)			
CROP	TARGET PESTS	RATE LBS AI/Acre (OZ./Acre)	REMARKS
ROOT, TUBER AND BULB VEGETABLES (continued)			
Turnips (FL, GA, IL, IN, OK, SC, TX, WA only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply within 1 day of harvest or feed crop refuse to livestock. Do not exceed 8 applications.
(TX, OK, IL, IN only)			Do not exceed 4 applications.
(SC, GA, FL, WA only)			Do not exceed 4 applications.
CEREAL GRAINS			
Field Corn/ Popcorn/ Field Corn Grain for Seed (Pre-emergent Use)**	Armyworm Cutworms	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply in the time period from 5 days prior to planting up to emergence of the crop.
Field Corn/ Popcorn/ Field Corn Grain for Seed (Foliar Use)**	Cutworms European Corn Borer Flea Beetle Corn Earworm*** Corn Rootworm Beetle*** Tobacco Budworm***	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Do not apply more than 0.6 lbs. a.i. per acre per season. Allow a minimum of 6 days between treatments. Do not apply within 30 days of harvest of grain or fodder (stover). Forage may be harvested on the day of application.
Stalk Borer			When treating for stalk borer, AMBUSH 25W must be applied when or shortly before the stalk borer larvae are moving into the corn from surrounding weeds and grasses. Mowing or burndown herbicide is suggested to initiate movement.**

(continued)

DIRECTIONS FOR APPLICATION (continued)			
CROP	TARGET PESTS	RATE LBS AI/Acre (OZ./Acre)	REMARKS
CEREAL GRAINS (continued)			
Sweet Corn	Corn Earworm European Corn Borer Fall Armyworm Corn Rootworm Adults Cutworms Aster Leafhopper Flea Beetle ^b	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 1.2 lbs. a.i. per acre per season. Do not apply within 1 day of harvest of ears or forage or livestock grazing. AMBUSH 25W may be mixed with LANIMATE [®] (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded.
Sweet Corn (Florida only)	Corn Earworm European Corn Borer Fall Armyworm Southern Armyworm Corn Rootworm Adults Cutworms Aster Leafhopper	0.1-0.25 (6.4-16.0)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 2.0 lbs. a.i. per acre per season. AMBUSH 25W may be mixed with LANIMATE [®] (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded.
TREE CROPS			
Apples	Apple Aphid Redbanded Leafroller Oblique Banded Leafroller Plum Curculio White Apple Leafhopper Spotted Tentiform Leafminer Tarnished Plant Bug	0.1-0.4 (6.4-25.6)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Do not apply after petal fall. Do not apply more than 0.6 lbs. a.i. per acre per season.

(continued)

DIRECTIONS FOR APPLICATION (continued)			
CROP	TARGET PESTS	RATE LBS AI/Acre (OZ./Acre)	REMARKS
TREE CROPS (continued)			
Avocados	Thrips Leafhoppers Mird Bugs Lepidopterous larvae	0.2 (12.8)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7 to 10 day intervals as needed to provide control. Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Cherries	Lesser Peachtree Borer Rose Chafer Green Fruitworm Redbanded Leafroller Plum Curculio Tarnished Plant Bug	0.1-0.2 (6.4-12.8)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Apply when insects appear. In the East, do not exceed 6 applications per season, with no more than 4 applications after petal fall. In the West, do not apply more than 4 applications per season, with no more than 3 applications after petal fall. Do not apply within 3 days of harvest. Do not graze livestock in treated orchards. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Filberts	Filbertworm Oblique Banded Leafroller	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 lbs. a.i. per acre per season. Do not apply within 14 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.

(continued)

RESTRICTED USE PESTICIDE**Due to Toxicity to Fish and Aquatic Organisms**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**ACTIVE INGREDIENT:****Permethrin**

(3-Phenoxyphenyl)methyl(±) cis, trans-3-(2,2-dichloroethyl)-
2,2-dimethylcyclopropanecarboxylate* 25.0%

INERT INGREDIENTS: 75.0%

TOTAL 100.0%

* Cis/trans ratio: Min. 35% (±) cis and max. 65% (±) trans.

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

EMERGENCY INFORMATION: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY: Transportation: CHEMTREC: 1-800-424-9300 Other: AMVAC: 1-323-264-3910

PRECAUTIONARY STATEMENTS. HAZARDS TO HUMANS AND DOMESTIC ANIMALS. WARNING: May cause eye irritation. Harmful if swallowed. Do not get in eyes, on skin or on clothing. **ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to fish and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

SEE ATTACHED BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

EPA Reg. No. 5481-502

EPA Est. No. 37429-GA-2



4100 E. Washington Blvd.
Los Angeles, CA 90023 U.S.A.

1-323-264-3910 • www.amvac-chemical.com

NET WEIGHT: 2 lbs

11483-3