



Bayer CropScience

Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina
 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

TEMIK® brand 15G Aldicarb Pesticide

For Use on Peanuts

EPA Reg. No. 264-330

EPA SLN. No. NC-810033

24(c) Supplemental Label

RESTRICTED USE PESTICIDE

For Retail Sale to and Use Only by Certified Applicators or Persons Under Their Direct Supervision and only for those Uses covered by the Certified Applicator's Certification

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH CAROLINA

DIRECTIONS FOR USE:

It is a violation of Federal and State Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

Dosage of TEMIK® 15G
Ounces Per 1000
Feet of Row
Pounds/
Acre
For 36" Row Spacing

Crop & Time of Application	Pests Controlled	Pounds/ Acre	Ounces Per 1000 Feet of Row	Recommended Application
PEANUTS: At time of planting and at peg initiation	INSECTS: Leafhopper, Mites (suppression)*	Apply 7 lbs. per acre at planting and at peg initiation	7.5 ozs. at planting 7.5 ozs. at peg initiation	For Control of insects and mites (suppression)*, apply granules in seed furrow and cover with soil. For control of nematodes, apply granules in a 6-12 inch band and work into soil or cover with soil to a depth of 2-4 inches. Plant seed in treated zone.
	Nematodes, Mites (suppression)*	Apply 10 lbs. per acre at planting and at peg initiation	11 ozs. at planting 11 ozs at peg initiation	At Peg initiation: For Control of insects, mites (suppression)* and nematodes, apply granules in a 12-18 inch band over the row and immediately incorporate into the soil with a suitable tilling device. In irrigated areas, follow pegging applications with irrigation within one week.

*Mite suppression can be expected from an at-planting treatment for control of leafhopper and a peg initiation treatment for control of nematodes. Treatments are ineffective on established infestations of mites.

Pre-Harvest and Grazing Use Information and Limitations

Peanuts

- Do not make more than one at-planting application and one post-emergence application per crop.
- Do not hog off treated fields.
- Do not feed hay or vines to livestock.
- Do not harvest within 90 days of last application.