



**Bayer CropScience**

**Bayer CropScience LP**  
**P.O. Box 12014**  
**2 T.W. Alexander Drive**  
**Research Triangle Park, North Carolina**  
**27709**  
**1-866-99BAYER (1-866-992-2937)**  
**<http://www.bayercropscienceus.com>**

**TEMIK® brand 15G Aldicarb Pesticide**

**For Use on Flue-Cured Tobacco**

**EPA Reg. No. 264-330**

**EPA SLN. No. NC-820008**

**24(c) Supplemental Label**

**RESTRICTED USE PESTICIDE**

For Retail Sale to and Use Only by Certified Applicators or Persons Under Their Direct Supervision and only for those Uses covered by the Certified Applicator's Certification

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH CAROLINA**

**DIRECTIONS FOR USE:**

It is a violation of Federal and State Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

**PLANT RESPONSE PRECAUTIONS:**

Under certain adverse weather conditions (e.g., high temperatures and/or dry weather) chlorosis and flecking of transplant leaves may occur. Symptoms may be more severe on greenhouse-grown transplants. Placement of transplant roots directly into a band narrower than recommended may result in phytotoxicity. If injury of this type is a concern, do not use this product.

Crop & Time of Application	Pests Controlled	Dosage of TEMIK® 15G		Recommended Application
		Pounds/ Acre (42-48 Inch Row Spacing)	Ounces Per 1000 Feet of Row	
<b>TOBACCO (Flue-Cured Only)</b> From one week before transplanting to immediately prior to transplanting. Do Not Apply After Transplanting	Aphids Flea Beetles* (Suppression)	10-14	14-20	Apply granules in a 6 to 12-inch band, and incorporate into or cover with soil to a depth of 2 to 6 inches when forming beds. Transplant into treated area.

## Alternate Application Method Recommended for Preformed Beds

### OFFSET SHANK APPLICATION PRIOR TO TRANSPLANTING:

Crop: TOBACCO (Flue-Cured Only)

Pest Controlled\*: Aphids

Pounds TEMIK® 15G/A: 10-14 (42 to 48-Inch Row Spacing)

Ounces TEMIK® 15G/1000 Row-Ft: 14-20 (42 to 48-Inch Row Spacing)

From one week before transplanting to immediately prior to transplanting, apply by shanking (one or two shanks per row) the material into the bed so that the granules are placed 4" to 5" to the side of the bed center and will be 5" below the final bed surface after transplanting (Figs. 1 and 2). If two shanks are used per row, 1/2 the total rate should be applied to each side of the row. Do not transplant roots directly into a shanked placement of TEMIK®, or phytotoxicity may occur. Do not apply after transplanting.

### Example of Equipment Setup and Granule Placement

Figure 1 - Shank Setup





Figure 2 - Deep Precision Placement



\*This application may provide early season flea beetle suppression.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

TEMIK is a registered trademark of Bayer.

08/09/04