

syngenta

ATRA

SPEND LESS. GROW MORE. LIVE BETTER.

**RESTRICTED USE PESTICIDE
(GROUND AND SURFACE WATER CONCERNS)**

**FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED
APPLICATOR'S OR PERSONS UNDER THEIR DIRECT
SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE
CERTIFIED APPLICATOR'S CERTIFICATION.**

**THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO
GROUND AND SURFACE WATER CONCERNS. USERS MUST READ
AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS
FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE
TO REACH GROUND AND SURFACE WATER.**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
OKLAHOMA**

AATREX® 4L

**EPA Reg. No. 100-497
EPA SLN No. OK-830003**

**For Control Of Annual Weeds, Using AAtrex 4L In Tank Mixtures,
During The Idle Season Following Winter Wheat Harvest In Early
Summer Until Planting Back To Winter Wheat That Fall.**

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA registered AAtrex 4L label and on the labels of herbicides tank mixed with AAtrex.

This labeling must be in the possession of the user at the time of pesticide application.

DO NOT contaminate or apply to ponds, lakes, or streams.

OK-830003

AAtrex is a broad spectrum herbicide which, in tank mixture with an EPA-registered herbicide for use in fallow, is effective for preemergence control of carpetweed, common lambsquarters, common purslane, kochia, mustards, Russian thistle, tumble and other pigweeds, wild lettuce, and suppression of volunteer wheat.

For best results, apply herbicide after straw has been uniformly distributed over the wheat stubble or soil surface.

Winter Wheat - After Harvest for Weed Control During Idle Season Until Fall Planting to Winter Wheat

Precautions: DO NOT apply on soils having a pH greater than 7. DO NOT apply in the Oklahoma Panhandle. DO NOT treat eroded hillsides, caliche and rocky outcroppings, or exposed calcareous subsoil. DO NOT treat soils with calcareous surface layers. DO NOT use on sands. Avoid spray overlap.

Apply AAtrex 4L at 0.5-1 pt./A in combination with an EPA-registered herbicide for use in a 4-month fallow, as soon as possible following wheat harvest.

Use the lower rates on coarse-textured soils and soils relatively low in organic matter; use the higher rates on fine-textured soils and soils relatively high in organic matter.

For control of established weeds, tank mix with paraquat at 1-2 pts./A + X-77® at 1-2 pts. per 100 gals. of spray solution, or stubble plow to control large established weeds. Refer to the herbicide labels for directions, weeds controlled, precautions, and limitations.

Apply in a minimum of 20 gals. of water per acre with ground equipment or in a minimum of 5 gals. of water per acre by air.

If weeds or volunteer wheat become a problem, a shallow cultivation or a postemergence application of paraquat + X-77 should be employed.

Note: DO NOT feed green forage or graze treated areas to livestock for six months following application. Treat only once during the same fallow period. DO NOT plant wheat until four months after application.

Chemigation: DO NOT apply these products through any type of irrigation system.

AAtrex® trademark of Syngenta
X-77® trademark of Loveland Industries, Inc.

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No.: OK-830003

Label Code:

Word: OKAAtrex4LWheatFallowSLNLbl9-17-03